



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

198. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

Three spheres of the same material, radii R , r , S , rest upon a horizontal plane, touching each other. Find the radius of a sphere of the same material as the others which, being placed upon the other three spheres, will just prevent the latter from separating, the coefficient of friction between the spheres being μ , and between the spheres and the table being μ' .

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let x =radius of top sphere center Q , R =radius of bottom sphere center P . The forces acting on the sphere radius R are its own weight W , the friction F , the friction F' , and the normal reaction between the two spheres R . Let nW_1 be the part of the weight of the sphere radius x supported by the sphere radius R . $\angle DPC=\beta$.

$$\text{Now } FHC=RDC. \quad F=R\left(\frac{DC}{HC}\right)=R\tan\frac{1}{2}\beta. \quad \therefore F=\mu R \text{ or } \mu=\tan\frac{1}{2}\beta.$$

$$F'=\mu'[W+nW_1]=R\sin\beta-F\cos\beta=R(\sin\beta-\tan\frac{1}{2}\beta\cos\beta)=R\tan\frac{1}{2}\beta.$$

$$\therefore R=\frac{\mu'[W+nW_1]}{\tan\frac{1}{2}\beta}=\frac{\mu'}{\mu}[W+nW_1].$$

$$\text{Resolving vertically, } nW_1=R\cos\beta+F\sin\beta.$$

$$\therefore nW_1=R[\cos\beta+\tan\frac{1}{2}\beta\sin\beta]=R. \quad \therefore nW_1=\frac{\mu'}{\mu}[W+nW_1].$$

$$\text{Let } \delta=\text{density of each sphere} \quad \therefore W=\delta g \times \frac{4}{3}\pi R^3, \quad W_1=\delta g \times \frac{4}{3}\pi x^3.$$

$$\therefore nx^3=\frac{\mu'}{\mu}[R^3+nx^3]. \quad \therefore n=\frac{\mu'R^3}{[\mu-\mu']x^3}.$$

Let mW_1 be the part of the upper sphere supported by sphere radius r , and pW_1 the part supported by sphere radius S . Then

$$m=\frac{\mu'r^3}{[\mu-\mu']x^3}, \quad p=\frac{\mu'S^3}{[\mu-\mu']x^3}.$$

But $n+m+p=1$.

$$\therefore \frac{\mu'[R^3+r^3+S^3]}{\mu-\mu'}=x^3, \text{ and } \therefore x=\sqrt[3]{\frac{\mu'[R^3+r^3+S^3]}{\mu-\mu'}}.$$

199. Proposed by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

A sphere of water, radius $\frac{1}{49}$, the earth's radius, is brought together by mutual attractions of particles from a state of infinite diffusion. Find its temperature owing to the amount of work done by these forces.

Solution by W. D. LAMBERT, Washington, D. C.

Let r be the variable radius of the sphere, while forming; a the final radius $= \frac{6.370 \times 10^8}{49}$ centimeters; ρ the density of water $= 1$; k the gravitation constant $= 6.665 \times 10^{-8}$ dynes; J the mechanical equivalent of heat $= 4.184 \times 10^7$ ergs for the centigrade gram-calorie. It is a little easier to conceive the sphere as pulled asunder against its own attraction, and the amount of work will be the same. Suppose the sphere made up of layers, each of thickness dr , and that the sphere has been reduced to radius r . The mass of a layer is $4\pi\rho r^2 dr$. The attraction between this mass and the remaining sphere is $\frac{k \times \frac{4}{3}\pi\rho r^3 \times 4\pi\rho r^2 dr}{x^2}$, where x denotes the distance of the layer (supposed to be scattered symmetrically) from the center of the sphere. The work done in removing the layer from the surface of the sphere in question to infinity is $\frac{1}{3}\pi^2 k \rho^2 r^5 dr \int_r^\infty \frac{dx}{x^2} = \frac{1}{3}\pi^2 k \rho^2 r^4$.

The total work done in removing all layers is

$$\frac{1}{3}\pi^2 k \rho^2 \int_0^a r^4 dr = \frac{1}{15}\pi^2 k \rho^2 a^5.$$

Dividing this quantity by the mass and by J we get for the temperature $\frac{4}{5} \frac{\pi k \rho a^2}{J}$. For substances other than water this result should be multiplied by the specific heat of the substance. Using the numerical values previously given, we get for the temperature $0^\circ.677$ centigrade.

Also solved by G. B. M. ZERR, whose result is 0.656. This difference of result is due to the different values assumed for the constants entering into the solution.

AVERAGE AND PROBABILITY.

183. Proposed by J. EDWARD SANDERS, Reinersville, Ohio.

A point within a given triangle is joined to each of the corners. What is the average of the sum of the lengths of these three lines?

I. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let ABC be the given triangle, P the random point, A the vertex, BC the base of the triangle, AD the altitude. Through P draw QR , parallel to BC cutting AD in F . Let $AD=p$, $BD=e$, $DC=d$, $AF=x$, $FP=y$. Then $AP=\sqrt{x^2+y^2}$. The limits of x are 0 and p ; of y , $-QF=ex/p$ and $+FR=dx/p$. Let M =average length of AP , Δ =average length of the sum.

$$\therefore M = \int_0^p \int_{-ex/p}^{dx/p} \sqrt{x^2+y^2} dx dy / \int_0^p \int_{-ex/p}^{dx/p} dx dy$$